

# Bibliographie

## Laryngectomie totale : Conséquences sur l'alimentation Déglutition, Olfaction, Gustation

### Ouvrages

Kerlan, M. (2016). *Éthique en orthophonie: Le sens de la clinique*. De Boeck Supérieur.

Ward, E. C., & van As-Brooks, C. J. (Eds.). (2014). *Head and neck cancer: treatment, rehabilitation, and outcomes*. Plural Publishing.

Woisard-Bassols, V., & Puech, M. (2011). *La réhabilitation de la déglutition chez l'adulte: le point sur la prise en charge fonctionnelle*. Groupe de Boeck.

### Référentiels

Institut National du Cancer (2019) « Les traitements des cancers des voies aérodigestives supérieures », *Guides de référence Cancer info*.

Haute Autorité de Santé (2010) « La prise en charge des cancers des voies aéro-digestives supérieures », *HAS - Guide patient*.

Recommandations SFORL [www.orlfrance.org/download.php?id=128](http://www.orlfrance.org/download.php?id=128)

Référentiels inter-régionaux de l'AFSOS, disponibles sur [www.afsos.org/-Referentiels-Nationaux-.html](http://www.afsos.org/-Referentiels-Nationaux-.html)

### Publications

#### Olfaction - Gustation

Hilgers, F. J., van Dam, F. S., Keyzers, S., Koster, M. N., van As, C. J., & Muller, M. J. (2000). Rehabilitation of olfaction after laryngectomy by means of a nasal airflow-inducing maneuver: the polite yawning technique. *Archives of Otolaryngology-Head & Neck Surgery*, 126(6), 726-732.

Hummel, T., Kobal, G., Gudziol, H., & Mackay-Sim, A. J. E. A. (2007). Normative data for the "Sniffin'Sticks" including tests of odor identification, odor discrimination, and olfactory thresholds: an upgrade based on a group of more than 3,000 subjects. *European Archives of Oto-Rhino-Laryngology*, 264(3), 237-243.

Kacha, S., Guillemin, F., & Jankowski, R. (2012). Development and validity of the DyNaChron questionnaire for chronic nasal dysfunction. *European Archives of Oto-Rhino-Laryngology*, 269(1), 143-153.

Risberg-Berlin, B., Rydén, A., Möller, R. Y., & Finizia, C. (2009). Effects of total laryngectomy on olfactory function, health-related quality of life, and communication: a 3-year follow-up study. *BMC Ear, Nose and Throat Disorders*, 9(1), 1-9.

Risberg-Berlin, B., Ylitalo, R., & Finizia, C. (2006). Screening and Rehabilitation of Olfaction After Total Laryngectomy in Swedish Patients: Results From an Intervention Study Using the Nasal Airflow-Inducing Maneuver. *Archives of Otolaryngology-Head & Neck Surgery*, 132(3), 301-306.

Soufflet Charlotte. (2020). *Mise en place d'une approche de réhabilitation olfactive pour la prise en charge des patients laryngectomisés totaux*. Nantes: Université de Nantes.

Ward, E., Coleman, A., Van As-Brooks, C., & Kerle, S. (2010). Rehabilitation of olfaction post-laryngectomy: a randomised control trial comparing clinician assisted versus a home practice approach. *Clinical Otolaryngology*, 35(1), 39-45.

## Déglutition après laryngectomie totale

De Casso, C., Slevin, N. J., & Homer, J. J. (2008). The impact of radiotherapy on swallowing and speech in patients who undergo total laryngectomy. *Otolaryngology–Head and Neck Surgery*, 139(6), 792-797.

Heijnen, B. J., Speyer, R., Kertscher, B., Cordier, R., Koetsenruijter, K. W. J., Swan, K., & Bogaardt, H. (2016). Dysphagia, speech, voice, and trismus following radiotherapy and/or chemotherapy in patients with head and neck carcinoma: review of the literature. *BioMed research international*, 2016. Hilgers, F. J., & Ackerstaff, A. H. (2000). Comprehensive rehabilitation after total laryngectomy is more than voice alone. *Folia phoniatrica et logopaedica*, 52(1-3), 65-73.

Hutcheson, K. A., Alvarez, C. P., Barringer, D. A., Kupferman, M. E., Lapine, P. R., & Lewin, J. S. (2012). Outcomes of elective total laryngectomy for laryngopharyngeal dysfunction in disease-free head and neck cancer survivors. *Otolaryngology--Head and Neck Surgery*, 146(4), 585-590.

Lazarus, C., Logemann, J. A., Shi, G., Kahrilas, P., Pelzer, H., & Kleinjan, K. (2002). Does laryngectomy improve swallowing after chemoradiotherapy?: A case study. *Archives of Otolaryngology–Head & Neck Surgery*, 128(1), 54-57.

Lennie, T. A., Christman, S. K., & Jadack, R. A. (2001). Educational needs and altered eating habits following a total laryngectomy. *Oncology Nursing Forum* (Vol. 28, No. 4).

Maclean, J., Cotton, S., & Perry, A. (2009). Dysphagia following a total laryngectomy: the effect on quality of life, functioning, and psychological well-being. *Dysphagia*, 24(3), 314-321.

Maclean, J., Szczesniak, M., Cotton, S., Cook, I., & Perry, A. (2011). Impact of a laryngectomy and surgical closure technique on swallow biomechanics and dysphagia severity. *Otolaryngology--Head and Neck Surgery*, 144(1), 21-28.

McConnel, F. M., Mendelsohn, M. S., & Logemann, J. A. (1986). Examination of swallowing after total laryngectomy using manofluorography. *Head & neck surgery*, 9(1), 3-12.

Mittal, B. B., Pauloski, B. R., Haraf, D. J., Pelzer, H. J., Argiris, A., Vokes, E. E., ... & Logemann, J. A. (2003). Swallowing dysfunction—preventative and rehabilitation strategies in patients with head-and-neck cancers treated with surgery, radiotherapy, and chemotherapy: a critical review. *International journal of radiation oncology, biology, physics*, 57(5), 1219-1230.

Organisation Mondiale de la Santé. (2001). *Classification internationale du fonctionnement, du handicap et de la santé: Cif*. Genève: Organisation mondiale de la Santé.

Pauloski, B. R., Logemann, J. A., Blom, E. D., & Hamaker, R. C. (1995). Functional outcome after surgery for prevention of pharyngospasms in tracheoesophageal speakers. Part II: Swallow characteristics. *The Laryngoscope*, 105(10), 1104-1110. Ward, E. C., Bishop, B., Frisby, J., & Stevens, M. B. B. S. (2002). Swallowing outcomes following laryngectomy and pharyngolaryngectomy. *Archives of Otolaryngology–Head & Neck Surgery*, 128(2), 181-186.

White, H. N., Golden, B., Sweeny, L., Carroll, W. R., Magnuson, J. S., & Rosenthal, E. L. (2012). Assessment and incidence of salivary leak following laryngectomy. *The Laryngoscope*, 122(8), 1796-1799.

## Questionnaires d'auto-évaluation

Audag, N., Goubau, C., Danse, E., Vandervelde, L., Liistro, G., Toussaint, M., & Reyckler, G. (2019). Validation and reliability of the French version of the Sydney swallow questionnaire. *Dysphagia*, 34(4), 556-566.

Balaguer, M., Percodani, J., & Woisard, V. (2017). The Carcinologic Handicap Index (CHI): A disability self-assessment questionnaire for head and neck cancer patients. *European annals of otorhinolaryngology, head and neck diseases*, 134(6), 399-403.

Crestani, S., Moerman, M., Woisard, V. (2011). The "Deglutition Handicap Index" a self-administrated dysphagia-specific quality of life questionnaire: sensibility to change. *Rev Laryngol Otol Rhinol*, 132(1), 3-7.

Fayers, P., Bottomley, A. E. O. R. T. C., & EORTC Quality of Life Group. (2002). Quality of life research within the EORTC—the EORTC QLQ-C30. *European Journal of Cancer*, 38, 125-133.

Govender, R., Lee, M. T., Davies, T. C., Twinn, C. E., Katsoulis, K. L., Payten, C. L., ... & Drinnan, M. (2012). Development and preliminary validation of a patient-reported outcome measure for swallowing after total laryngectomy (SOAL questionnaire). *Clinical Otolaryngology*, 37(6), 452-459.

Kacha, S., Guillemin, F., & Jankowski, R. (2012). Development and validity of the DyNaChron questionnaire for chronic nasal dysfunction. *European Archives of Oto-Rhino-Laryngology*, 269(1), 143-153.

Lechien, J. R., Cavelier, G., Thill, M. P., Bousard, L., Blecic, S., Vanderwegen, J., ... & Dequanter, D. (2020). Validity and reliability of a French version of MD Anderson Dysphagia Inventory. *European Archives of Oto-Rhino-Laryngology*, 277, 3111-3119.

Singer, S., Araújo, C., Arraras, J. I., Baumann, I., Boehm, A., Brokstad Herlofson, B., ... & EORTC Quality of Life and the EORTC Head and Neck Cancer Groups. (2015). Measuring quality of life in patients with head and neck cancer: Update of the EORTC QLQ-H&N Module, Phase III. *Head & neck*, 37(9), 1358-1367.

